

Science News*from research organizations*

SARS-CoV-2 spike proteins disrupt the blood-brain barrier, new research shows

Potentially raising risk of neurological damage in COVID-19 patients

Date: October 29, 2020

Source: Temple University Health System

Summary: New research shows that the spike proteins that extrude from SARS-CoV-2 promote inflammatory responses on the endothelial cells that form the blood-brain barrier. The study shows that SARS-CoV-2 spike proteins can cause this barrier to become 'leaky,' potentially disrupting the delicate neural networks within the brain.

Share: [!\[\]\(d3102649f02e825ddb76dc3de0190154_img.jpg\)](#) [!\[\]\(55ca3a38dbb940110628e54e3ea7505d_img.jpg\)](#) [!\[\]\(1ad7b9dfa1e10e48660df5dd18a1b20c_img.jpg\)](#) [!\[\]\(7b7f78f3b14c2b344e3d1b2a79a760c9_img.jpg\)](#) [!\[\]\(ea85bb5cc605bcd0e585ecc5f09976f5_img.jpg\)](#)

FULL STORY

Like a key, SARS-CoV-2 -- the virus that causes coronavirus disease 2019 (COVID-19) -- attaches to specific molecules on the host cell surface, opening gateways into the cell interior. Viral entry into host cells triggers a prodigious immune response. Much of this battle is waged within the lungs, which explains why many patients hospitalized with COVID-19 have severe respiratory symptoms.

Respiratory symptoms, however, are only part of the story. Increasing evidence points toward blood vessel inflammation as having a crucial impact on the severity of COVID-19. In addition, anywhere from 30 to 80 percent of patients experience neurological symptoms, including dizziness, headache, nausea, and loss of concentration. These symptoms suggest that SARS-CoV-2 also affects cells of the central nervous system.

While there is no evidence yet that the virus invades the brain, new work by scientists at the Lewis Katz School of Medicine at Temple University shows that the spike proteins that extrude from SARS-CoV-2 promote inflammatory responses on the endothelial cells that form the blood-brain barrier. The study, published in the December print issue of the journal *Neurobiology of Disease*, is the first to show that SARS-CoV-2 spike proteins can cause this barrier to become "leaky," potentially disrupting the delicate neural networks within the brain.

"Previous studies have shown that SARS-CoV-2 infects host cells by using its spike proteins to bind to the angiotensin converting enzyme 2 (ACE2) on the host cell surface," explained Servio H. Ramirez, PhD, Professor of Pathology and Laboratory Medicine at the Lewis Katz School of Medicine at Temple University and principal investigator on the new study.

ACE2 is expressed on endothelial cells, which form the inner lining of blood vessels, and serves a central role in mediating different functions of the cardiovascular system. According to Dr. Ramirez, "since ACE2 is a major binding target for SARS-CoV-2 in the lungs and vasculature of other organs in the body, tissues that are behind the vasculature, that receive blood from affected vessels, are at risk of damage from the virus."

It has been unclear, however, whether ACE2 is also present in the brain vasculature or whether its expression changes in health conditions that worsen COVID-19, such as high blood pressure (hypertension). To find out, the team began by examining postmortem human brain tissue for vascular ACE2 expression, using tissues from individuals without underlying health conditions and from individuals in whom hypertension and dementia had been established. Analyses showed that ACE2 is in fact expressed throughout blood vessels in the frontal cortex of the brain and is significantly increased in the brain vasculature of persons with a history of hypertension or dementia.

The researchers then investigated the effects of the SARS-CoV-2 spike protein on brain endothelial cells in cell culture models. Introduction of the spike protein, particularly a portion designated subunit 1, produced substantial changes in endothelial barrier function that led to declines in barrier integrity. The researchers also uncovered evidence that subunit 2 of the SARS-CoV-2 spike protein can directly impact blood-brain barrier function. "This is of importance because unlike subunit 1, subunit 2 of the spike protein doesn't bind to ACE2, meaning that a breach to the blood-brain barrier could occur in a manner that is independent of ACE2," explained postdoctoral fellow and first author on the new report Tetyana P. Buzhdygan, PhD.

Dr. Ramirez's team further investigated the effects of SARS-CoV-2 spike proteins on tissue-engineered microfluidic constructs designed to mimic a human brain capillary. "The tissue-engineered microfluidic models allow recapitulation of the 3D cyto-architecture and mechanical forces caused by fluid movement, which the vasculature is continuously exposed to," said Allison M. Andrews, PhD, Assistant Professor in the Department of Pathology & Laboratory Medicine at LKSOM and a co-author on the report. Experiments showed that binding of spike protein subunit 1 increased barrier permeability in the engineered vessel-like constructs.

"Our findings support the implication that SARS-CoV-2, or its shed spike proteins circulating in the blood stream, could cause destabilization of the blood-brain barrier in key brain regions," Dr. Ramirez said. "Altered function of this barrier, which normally keeps harmful agents out of the brain, greatly increases the possibility of neuroinvasion by this pathogen, offering an explanation for the neurological manifestations experienced by COVID-19 patients."

The long-lasting effects of altered blood-brain barrier function in the presence of SARS-CoV-2 are unknown. Moreover, as Dr. Buzhdygan explained, "the brain vasculature is extremely branched, so even a small amount of neuroinflammation can be very damaging." Based on the team's observations of ACE2 expression in the brain, this neurological damage could be extensive in COVID-19 patients with pre-existing health conditions in which the vasculature has already suffered some amount of injury.

It also remains unknown whether the virus can actually get inside neurons or glial cells that lie beyond the barrier. "The viral genome has not been found yet in the specific cell types of the brain," Dr. Ramirez noted. "The next steps in our work are to look for genomic viral copies in different parts of the brain using autopsy material from COVID-19 cases and to investigate the pathogen's ability to neuroinvasion using different cell culture and tissue-engineered constructs."

Other key collaborators on the project include Dr. Raghava Potula, Department of Pathology and Laboratory Medicine at LKSOM, and Dr. Peter A. Galie, Department of Biomedical Engineering, Rowan University, New Jersey.

Story Source:

Materials provided by **Temple University Health System**. *Note: Content may be edited for style and length.*

Journal Reference:

1. Tetyana P. Buzhdygan, Brandon J. DeOre, Abigail Baldwin-Leclair, Trent A. Bullock, Hannah M. McGary, Jana A. Khan, Roshanak Razmpour, Jonathan F. Hale, Peter A. Galie, Raghava Potula, Allison M. Andrews, Servio H. Ramirez. **The SARS-CoV-2 spike protein alters barrier function in 2D static and 3D microfluidic in-vitro models of the human blood-brain barrier.** *Neurobiology of Disease*, 2020; 146: 105131 DOI: 10.1016/j.nbd.2020.105131
-

Cite This Page:

MLA	APA	Chicago
-----	-----	---------

Temple University Health System. "SARS-CoV-2 spike proteins disrupt the blood-brain barrier, new research shows: Potentially raising risk of neurological damage in COVID-19 patients." ScienceDaily. ScienceDaily, 29 October 2020. <www.sciencedaily.com/releases/2020/10/2021029141941.htm>.

RELATED STORIES

SARS-CoV-2 Infections May Trigger Antibody Responses Against Multiple Virus Proteins

June 21, 2021 — All coronaviruses produce four primary structural proteins and multiple nonstructural proteins. However, the majority of antibody-based SARS-CoV-2 research has focused on the spike and nucleocapsid ...

Scientists Identify Virus-Cell Interaction That May Explain COVID-19's High Infection Rate

Mar. 29, 2021 — Researchers quantify the specific interaction between the spike protein of SARS-CoV-2 -- the virus that causes COVID-19 -- with the ACE2 receptors in human cells that may partially explain its high ...

T Cells Can Mount Attacks Against Many SARS-CoV-2 Targets -- Even on New Virus Variant

Jan. 27, 2021 — A new study suggests that T cells try to fight SARS-CoV-2 by targeting a broad range of sites on the virus -- beyond the key sites on the virus's spike protein. By attacking the virus from many ...

COVID-19 Virus Enters the Brain, Research Strongly Suggests

Dec. 17, 2020 — The SARS-CoV-2 virus, like many viruses before it, is bad news for the brain. In a new study, researchers found that the spike protein, often depicted as the red arms of the virus, can cross the ...

FROM AROUND THE WEB

ScienceDaily shares links with sites in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Small intestine cells at risk for COVID-19 infection in IBD patients

Healio

Spike proteins provide 'plausible explanation' for neurological complications of COVID-19

Healio

Genome-wide CRISPR activation screen identifies candidate receptors for SARS-CoV-2 entry

Shiyou Zhu et al., SCIENCE CHINA Life Sciences, 2021

A novel coronavirus—from basic research to clinic

Dan LUO et al., SCIENTIA SINICA Vitae, 2021

Realization of humoral immunity against SARS-CoV-2 infections

Lin et al., Fundamental Research, 2021

'We have more to learn': Data inconclusive on COVID-19 driving form of type 1 diabetes

Healio

Sysmex Obtains Japan Marketing Approval to Distribute BGI Genomics Coronavirus Kit | Genomeweb

staff reporter, GenomeWeb

In Brief This Week: Illumina, Guardant Health, PerkinElmer, and More | Genomeweb

staff reporter, GenomeWeb

Powered by **TREND MD**

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). [Learn more](#)

Yes

No

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

✉ Email Newsletters

📡 RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

f Facebook

🐦 Twitter

in LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

💬 Leave Feedback

📞 Contact Us

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2021 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners.

Content on this website is for information only. It is not intended to provide medical or other professional advice.

Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— GDPR: Privacy Settings —